mlbio 酶联生物
Good elisakit producers
订购热线: 4008-898-798

Anti-DYNC1LI2 antibody

Cat. No.	ml263749
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description	Anti-DYNC1LI2 rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Synthetic peptide of human DYNC1LI2
Reactivity	Human, Mouse, Rat
Content	0.9 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	DYNC1LI2
Full name	dynein cytoplasmic 1 light intermediate chain 2

mlbio 酶联生物 Good elisakit producers

mlbio 酶联生物
Good elisakit producers
订购热线: 4008-898-798

Synonyms	LIC2; DNCLI2
Swissprot	O43237

Target Background

Acts as one of several non-catalytic accessory components of the cytoplasmic dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. May play a role in binding dynein to membranous organelles or chromosomes.

mlbio 酶联生物
Good elisakit producers
订购热线: 4008-898-798

Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human liver cancer

Recommended dilution: 20-100





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml263749(DYNC1LI2 Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: ×200)

Western blotting

Predicted band size:54 kDa

Positive control:Hela and PC-3 cell, Mouse brain tissue, SKOV3 cell, Human fetal brain tissue lysates

Recommended dilution: 200-1000

mlbio 酶联生物
Good elisakit producers

mlbio 酶联生物
Good elisakit producers
订购热线: 4008-898-798

Gel: 8%SDS-PAGE

Lysate: 40 μg

Lane 1-5: Hela and PC-3 cell, Mouse brain tissue, SKOV3 cell, Human fetal brain tissue lysates

Primary antibody: ml263749(DYNC1LI2 Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 3 minutes



ELISA

Recommended dilution: 5000-10000

联系电话：4008-898-798，021-61725725

联系QQ：2881505695，2881505696

邮箱：mlbio_cn@yeah.net

网址：www.mlbio.cn