

订购热线: 4008-898-798

Anti-HEBP1 antibody

Cat. No.	ml264155
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Product overview	
Description	Anti-HEBP1 rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Synthetic peptide of human HEBP1
Reactivity	Human
Content	1.08 mg/ml
Host species	Rabbit
lg class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification
	Antigen affinity purification
Target information	oroca
Symbol	HEBP1
Full name	heme binding protein 1
(-,00



Synonyms HBP; HEBP

Swissprot Q9NRV9

Target Background

The full-length protein encoded by this gene is an intracellular tetrapyrrole-binding protein. This protein includes a natural chemoattractant peptide of 21 amino acids at the N-terminus, which is a natural ligand for formyl peptide receptor-like receptor 2 (FPRL2) and promotes calcium mobilization and chemotaxis in monocytes and dendritic cells.



订购热线: 4008-898-798


Applications


Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human tonsil

Recommended dilution: 40-200





Good elisa Nit producer

The image on the left is immunohistochemistry of paraffinembedded Human tonsil tissue using ml264155(HEBP1 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)

Western blotting


Predicted band size:21 kDa


Positive control:Human kidney tissue and 293T cell lysates

Recommended dilution: 500-2000

Predicted cell location: Cytoplasm Positive control: Human prostate cancer

Recommended dilution: 40-200





The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using ml264155(HEBP1 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)



Gel: 12%SDS-PAGE	kDa	1	2
Lysate: 40 µg	95— 72—	_	
Lane 1-2: Human kidney tissue and 293T cell lysates	55—		
Primary antibody: ml264155(HEBP1 Antibody) at dilution 1/450	36— 28—		
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution		-	-
	17 <u></u> 10 <u></u>		
Exposure time: 30 seconds			

ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn