

磷酸化蛋白激酶Cn抗体

产品货号: mlR7382

英文名称: phospho-PKC eta (Thr656)

中文名称: 磷酸化蛋白激酶 Cn 抗体

别名: PKC eta (phospho T656); p-PKC eta (phospho T656); KPCL_HUMAN; MGC 5363; MGC 5363; MGC 5363; MGC26269; MGC5363; nPKC eta; nPKC-eta; PKC h; PKC L; PKC-L; PKCh; PKCL; PRKCH; PRKCL; Protein kinase C eta; Protein kinase C eta type.

产品类型: 磷酸化抗体

研究领域: 细胞生物 信号转导 激酶和磷酸酶

- 抗体来源: Rabbit
- 克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat,

产品应用: WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 80kDa

细胞定位: 细胞浆 细胞膜

性 状: Lyophilized or Liquid



浓 度: 1mg/ml

免疫原: KLH conjugated synthesised phosphopeptide derived from human PKC eta around the phosphorylation site of Thr656:VL(p-T)PI

亚型: lgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍: Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipids-dependent protein kinase. It is predominantly expressed in epithelial tissues and has been shown to reside specifically in the cell nucleus. This protein kinase can regulate keratinocyte differentiation by activating the MAP kinase MAPK13 (p38delta)-activated protein kinase cascade that targets CCAAT/enhancer-binding protein alpha (CEBPA). It is also found to mediate the transcription activation of the transglutaminase 1 (TGM1) gene. [provided by RefSeq, Jul 2008]

Function:

This is calcium-independent, phospholipid-dependent, serine- and threonine-specific enzyme.

PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters.



Tissue Specificity:

Most abundant in lung, less in heart and skin.

DISEASE:

Defects in PRKCH may be a cause of susceptibility to ischemic stroke (ISCHSTR) [MIM:601367]; also known as cerebrovascular accident or cerebral infarction. A stroke is an acute neurologic event leading to death of neural tissue of the brain and resulting in loss of motor, sensory and/or cognitive function. Ischemic strokes, resulting from vascular occlusion, is considered to be a highly complex disease consisting of a group of heterogeneous disorders with multiple genetic and environmental risk factors.

Similarity:

Belongs to the protein kinase superfamily.

AGC Ser/Thr protein kinase family.

PKC subfamily.

Contains 1 AGC-kinase C-terminal domain.

Contains 1 C2 domain.

Contains 2 phorbol-ester/DAG-type zinc fingers.

Contains 1 protein kinase domain.

SWISS:

P24723

Gene ID:

5583



Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.