

## 白细胞介素 28 受体 $\alpha$ 抗体

产品货号： mlR18148

英文名称： IL28 Receptor alpha

中文名称： 白细胞介素 28 受体  $\alpha$  抗体

别名： Class II cytokine receptor CRF2/12; CRF2-12; CRF2/12; Cytokine receptor class-II member 12; Cytokine receptor family 2 member 12; IFN lambda R1; IFN-lambda receptor 1; IFNLR; IFNLR1; IL-28 receptor subunit alpha; il-28r1; IL28R alpha; IL28RA; Interferon lambda receptor; Interferon lambda receptor 1; interleukin 28 alpha receptor; interleukin 28 receptor A; Interleukin 28 receptor alpha chain; interleukin 28 receptor, alpha (interferon, lambda receptor); Interleukin 28 receptor, alpha; interleukin or cytokine receptor 2; Interleukin-28 receptor subunit alpha; LICR2; Likely interleukin or cytokine receptor 2.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 56kDa

细胞定位： 细胞膜

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human IL28 Receptor alpha:151-250/520  
<Extracellular>

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** The protein encoded by this gene belongs to the class II cytokine receptor family. This protein forms a receptor complex with interleukine 10 receptor, beta (IL10RB). The receptor complex has been shown to interact with three closely related cytokines, including interleukin 28A (IL28A), interleukin 28B (IL28B), and interleukin 29 (IL29). The expression of all three cytokines can be induced by viral infection. The cells overexpressing this protein have been found to have enhanced responses to IL28A and IL29, but decreased response to IL28B. Three alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

**Function:**

IL28RA is a class II cytokine receptor that interacts with IL10RB to bind IL28A, IL28B, and IL29, which are related to type I interferons. These ligands play a critical role in response to microbial challenge and activate the JAK/STAT signaling system. It heterodimers with IL10RB. There are 4 isoforms as result of alternative splicing. It is widely expressed. IL28RA belongs to the type II cytokine receptor family, and contains 1 fibronectin type III domain.

**Subunit:**

Heterodimer with IL10RB.

**Subcellular Location:**

Membrane; Single-pass type I membrane protein

**Tissue Specificity:**

Widely expressed.

**Similarity:**

Belongs to the type II cytokine receptor family.

Contains 1 fibronectin type-III domain.

**SWISS:**

Q8IU57

**Gene ID:**

163702

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.