

黑色素瘤相关抗原 D2 抗体

产品货号： mlR18625

英文名称： MAGED2

中文名称： 黑色素瘤相关抗原 D2 抗体

别名： 11B6; 11B6; BCG 1; Breast cancer associated gene 1 protein; HCA10; HCA10; Hepatocellular carcinoma associated protein JCL 1; JCL 1; MAGE D; MAGE D2; MAGE D2 antigen; MAGE D2 antigen; Melanoma antigen family D 2; Melanoma antigen family D 2; Melanoma associated antigen D2; Melanoma associated antigen D2; MGC8386; MGC8386; Moderately similar to NECDIN [Mus musculus].

研究领域： 肿瘤 细胞生物

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 49kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human MAGED2:131-230/436

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : This gene is a member of the MAGED gene family. The MAGED genes are clustered on chromosome Xp11. This gene is located in Xp11.2, a hot spot for X-linked mental retardation (XLMR). This gene may also be involved in several types of cancer, including breast cancer and melanoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2013]

Subcellular Location:

Cytoplasmic

Tissue Specificity:

Widely expressed in normal tissues.

Similarity:

Belongs to the intermediate filament family.



SWISS:

Q9UNF1

Gene ID:

10916

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.